

July 13, 2010

filename bibh.wpd

Haeckel, ?. 1866. Generelle Morphologie der Organismen, vol. 2, p. lxxvii, 57, 77.

Hagadorn, J. W. 2002. Bear Gulch: an exceptional Upper Carboniferous Plattenkalk. pp. 167-183 in D. J. Bottjer, W. Etter, J. W. Hagadorn & C. M. Tang (eds.) Exceptional fossil preservation: a unique view on the evolution of marine life. Columbia Univ. Press, NY 403 pp.

Hagadorn, J. W., T. E. Whiteley and K. H. Nealson. 2001. 3D imaging of pyritized soft tissues in Paleozoic Konservat-Lagerstätten. GSA Annual Meeting Nov. 2001, Boston. Abstract. [Urasterella asperula] [Hunsrück Slate]

Hahn, Gerhard and Carsten Brauckmann. 1981. Ein neuer Ophiuren-Fund aus dem Kulm von Herborn (Asterozoa, Unter-Karbon III α , Hessen). Geologisches Jahrbuch Hessen, 109: 5-18, Wiesbaden. [source Pek & Prokop 1986] [Chattaster n.g. (Ophiurinae) includes C. dillensis n.sp. and Lapworthura hueffneri Schöndorf 1915] [mention of Ophiuroidea gen. et sp. indet E. Schmidt 1930, Xenura koboldi Schöndorf 1932, and Silesiaster longivertebralis Schwartzbach & Zimmermann 1936]

Hakes, W. G. 1976. Trace fossils and depositional environment of four clastic units, Upper Pennsylvanian megacyclothems, northeast Kansas. The University of Kansas Paleontological Contributions, Article 63. [Asteriacites pp. 20-21, Rock Lake Shale Member and Stull Shale Member]

Hakes, W. G. 1977. Trace fossils in Late Pennsylvanian cyclothems, Kansas. In T. P. Crimes & J. C. Harper (eds.), Trace fossils 2: Geol. Journal Spec. Issue 9, Liverpool, Seel House Press, pp. 209-22. [source Hakes 1985; Lawrence Shale Asteriacites quinquefolis] [p. 213, A.q. at locality 3a (measured section) SW $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 25, T.12S., R.19E.] [from Hakes 1985 Fig. 2 this locality is Douglas County]

Hakes, William G. 1985. Trace fossils from brackish-marine shales, Upper Pennsylvanian of Kansas, U.S.A. pp. 21-36 in H.A. Curran (ed.) Biogenic structures: their use in interpreting depositional environments. Society of Economic Paleontologists and Mineralogists Special Publication No. 35, 347 pp. [Asteriacites lumbricalis 12-15mm diameter in Rock Lake Shale member of Stanton Limestone, Lansing Group (Missourian stage)] [Asteriacites quinquefolis 15mm in diameter in "Lawrence Shale" member of Lawrence Formation, Douglas Group (Virgilian stage)]

Halfar, A. 1893. Die erste Asteride aus den palaozoischen Schichten des Harzes. Jahrb. k. preuss. geol. Landesanst. u. Bergakad., Berlin, 1892, vol. 13, pp. 186-199, pl. 10.

July 13, 2010

- Hall, James. 1847. Natural History of New York. Part 6, Paleontology, vol. 1, containing descriptions of the organic remains of the Lower Division of the New York System, 338 pp., 99 pls. [source Golden & Nitecki 1970] [p. 18 and Plate 4 fig. 11 Asterias? (sp. indet) but see Orbigny 1850 p. 22 = Asterias tenuiradiatus = Coelaster tenuiradiatus] [Asterias matutina Hall p. 91 and Pl. 29 fig. 5]
- Hall, James. 1852. Natural History of New York. Part 6, Palaeontology, vol. 2, containing descriptions of the organic remains of the Lower Middle Division of the New York System, 363 pp., 105 pls. [source Golden & Nitecki 1970] [p. 247]. [Palaeaster niagarensis Hall]
- Hall, James. 1859. Natural History of New York. Palaeontology, vol. 3, p. 134.
- Hall, James. 1861. Descriptions of new species of Crinoidea [from investigations of the Iowa Geological Survey.] Preliminary notice. 19 pp. Albany. [p. 18: Protaster? barrisi] [= Onychaster]
- Hall, James. 1868. Contributions to palaeontology. 12. Note on the genus Palaeaster and other fossil starfishes; with descriptions of some new species, and observations upon those previously described.--New York State Cabinet of Natural History Twentieth Report: 282-301; revised ed. 1868 = 1870:324-345 + pl. 9. [Also published in advance of Report in 8o, 24 pp., pl. 9, in Nov. and Dec. 1866 as noted under "Scientific Intelligence" in the American Journal of Science and Arts, second series, vol. XLIII, whole number XCIII, no. 129, p. 409.] [Palaeaster eucharis Hall]
- Hall, James. 1885. Note on the intimate relations of the Chemung group and the Waverly sandstone in northwestern Pennsylvania and southwestern New York. Proc. Am. Ass. Ad. Sci., vol. 33, pp. 416-419. [Paper presented at 33rd mtg in 1884; starfishes on p. 417; specimens in YPM.]
- Hall, Townsend M. 1867. On the relative distribution of fossils throughout the North Devon Series. Quart. Journ. Geol. Soc. Lond., vol. 23, pt. I [Proceedings of the Geological Society], pp. 371-381.
- Halpern, Jeanne W. 1972. Tracking down fossils at the Museum of Paleontology. Research News, Office of Research Administration, The University of Michigan, vol. XXII, no. 7/8, 14 pp. [Strataster ohioensis figured, anecdote of its discovery, cover & pp. 8, 9.]
- Hammann, W. and S. Schmincke. 1986. Depositional environment and systematics of a new ophiuroid, Taeniaster ibericus n. sp., from the Middle Ordovician of Spain.--Neus Jahrbuch für Geologie und Paläontologie Abhandlungen 173:47-74.

July 13, 2010

- Han, Nai-Ren; Zhang, Jia-Zhi; Zhang, Shi. 2004. A new species of Asterozoa from the Middle Devonian of Xiangzhou, Guangxi. *Acta Palaeontologica Sinica* 43(1), Jan 2004:142-146. [In Chinese with English summary] [ZR 2004] [Devonaster guangxiensis sp. nov., Guannxi Zhuang, Devonian]
- Hand, Greg. 1998. Unique collector Bill White dies. -- Dry Dredgers 9/98. Reprinted in Oct-Nov 1998 issue of MAPS DIGEST 21(7-8):3. [□.the crinoid and starfish pocket at Hueston Woods, is now deposited at Miami University]
- Hansen, T., D. L. Bruton and S. L. Jakobsen. 2005. Starfish from the Ordovician of the Oslo Region, Norway. *Norwegian Journal of Geology* 85(3):209-216. [Cnemidactis osloensis n. sp., Elnes Formation (Llanvirn); Stenaster obtusus, Furuberg Fm. (Caradoc)] [Arenig asteroid material from the Leningrad region (fide Rozhnov)] [C? macroadambulacralatas Blake & Guensburg 1993 is undoubted Cnemidactis]
- Hansman, Shaw and Pettyjohn. 1962. Supplement to the catalog of the type specimens of fossils in the University of Cincinnati Museum. University of Cincinnati, pp. iii + 131. [Protasterina fimbriata pp. 2, 102; Hexura weitzii p. 46.] [see Jell & Theron 1999 for Hexuraster replacement name]
- Häntzschel, W. 1975. Trace fossils and Problematica. In *Treatise on Invertebrate Paleontology, Part W, Miscellanea, Supplement 1*, C. Teichert (ed.). GSA and University of Kansas Press, 269 pp.
- Harland, W. B., et. al., eds. 1967. *The fossil record*. Geol. Soc. London. 828 pp. [source Sepkoski 1982]
- Harnack, Curt. 1959. Treasures from an ancient sea (Iowa). *Earth Science* (Chicago, Ill.) vol. 12, no. 5, pp. 153-156, 166, illus., reprinted Oct. 1959.
- Harper, D. A. T. 1981. The Late Ordovician Lady Burn Starfish Beds of Girvan. *Proceedings of the Geological Society of Glasgow* (Sessions 122/123) 1980-1981 (unpaginated separate). [burial occurred during sudden downslope movement of both sediment and fauna] [considerable downslope transport] [unstable submarine fan] [shelf and slope faunas in the beds]
- Harper, G. W. and R. S. Bassler. 1896. *Catalogue of the fossils of the Trenton and Cincinnati Periods, occurring in the vicinity of Cincinnati, Ohio*. Cincinnati, printed by The W. B. Carpenter Co., 810-314 Walnut St. pp. iv + 34. [Asteroids pp. 6-7, nos. 105-122; ophiuroids p. 7, nos. 123-127.] [Taeniaster elegans, T. fimbriatus, T. flexuosus, T. granuliferus, T. miamiensis.]
- Harper, John A. 1985. A new look at Eugasterella logani (Hall, 1868) (Stelleroidea:

July 13, 2010

Ophiuroidea) from the Middle Devonian of New York State.--Annals of Carnegie Museum 54:357-373.

Harper, John A., and Helen L. Delano. 2005. Reflections on a Devonian brittle star from the shore of Lake Erie. Pennsylvania Geology 35(3/4):2-9 [plus cover illustration; plus explanation of the cover] [encrinasterid, gen. et sp. indet] [weathered concretion from Northeast Shale; calcareous shale slab]

Harper, John A. and Robert W. Morris. 1978. A new encrinasterid ophiuroid from the Conemaugh Group (Pennsylvanian) of western Pennsylvania, and revision of the Encrinasteridae. Journal of Paleontology, vol. 52, no. 1, pp. 155-163 [Armathyraaster paradoxis n. g., n. sp.; Armathyraasterinae n. subfam.] [Pennsylvanian Brush Creek Shale, Glenshaw Formation; Punxsutawney, Jefferson County, Pennsylvania]

Hattin, D. E. 1959. An occurrence of Ophiuraster barrisi Miller. Jour. Paleont., vol. 33, pp. 1125-1126, 1 text-fig.

Hattin, D. E. 1967. Permian ophiuroids from Northern Oklahoma. Jour. Paleont., vol. 41, no. 2, pp. 489-492, 3 text-figs., 1 table.

Haude, Reimund 1982. Ophiuren (Echinodermata) aus dem Karbon des Rheinischen Schiefergebirges. Geologisches Jahrbuch Hessen 110:5-26, Weisbaden. [source Pek & Prokop 1986] [Pectinura horni n. gen., n. sp. (Klasmuridae), comet form with six arms; Protasteracanthion discussed] [Furcaster n. sp.? vertebrae figured/described; Chattaster assigned to Protasteridae, monotypic C. hueffneri (=C. dillensis)]

Haude, Reimund . 1992. Hertha Sieverts-Doreck. Jb. Ges. Naturkde. Württemberg 147: 349-356. [obituary and bibliography; author of some papers on Paleozoic Asterozoa]

Haude, Reimund . 1995. Echinodermen aus dem Unter-Devon der argentinischen Präkordillere. N. Jb. Geol. Pal., Abh., 197:37-86, 16 Abb. [Promopalaeaster? quadriserialis n.sp.; Marginaster yachelensis (Ruedemann), Argentinaster bodenbenderi, Furcaster separatus n.sp.]

Haude, Reimund . 1996. Fluchversuche verschütteter Holothurien in einer unterdevonischen Echinodermen-Taphozönose. Terra Nostra 96/6:48. [L. Devonian asteroids and ophiuroids also present]

Haude, Reimund . 1998. Die Ophiuren (Echinodermata) aus den Fundschichten Bodenbenders im Unter-Devon der argentinischen Präkordillere. Terra Nostra 98/5:65. [eight species. 1) Argentinaster bodenbenderi, 2) "Encrinaster" yachalensis, 3) Furcaster sp., 4) Urosoma? n.sp., 5) Eospondylus? n. sp., 6) "A-ura" n.sp. cf Ophiurinae, 7) "B-ura" n.sp. cf Protasteridae, 8) "C-ura" n.sp. cf Protasteridae]

July 13, 2010

- Haude, Reimund. 1999. Der-verzögerte-Ersatz eines Homonyms: Marginaster Haude 1995. [The delay in replacing a homonym.] N. Jb. Geol. Paläont. Mh. 1999(5):292-294. [replacement is Marginura. Advocates subfamily Euzonosomatinae for Euzonosoma and Marginura.]
- Haude, Reimund. 2004. Small skeletal elements of late Famennian echinoderms from the Refrath 1 Borehole [Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany]. Cour. Forsch.-Inst. Senckenberg 251:237-251. [ophiuroid ossicles identified, figured: Klasmuridae, ?Encrinasteridae, ?Protasteridae, Furcaster sp., Ophiurinae]
- Haude, Reimund. 2004. Morphology and palaeobiology of echinoderms in the Lower Devonian of the Argentine Precordillera. pp. 417-419 in T. Heinzeller & J. H. Nebelsick (eds.) Echinoderms: München. Taylor & Francis Group, London. xxvi + 633 pp. [includes paleobiologic placement of Argentinaster and an eospondylid] [9 species of ophiuroids, 2 species of asteroid, Talacasto Fm]
- Haude, Reimund & F. H. C. Hotchkiss. 2003. Observations on Aganaster gregarius and Stephanoura belgica (Ophiuroidea: Ophiolepididae). Program and Abstracts, 11th IEC, Munich. [see also Hotchkiss & Haude 2004]
- Haude, Reimund, and A. F. Sterren. 2007. Biostratigraphy and palaeobiology of echinoderms in the Lower Devonian of the Argentine Precordillera. Field Meeting of the IGCP 499 (IUGS/UNESCO), San Juan, Argentina May 14-22, 2007. Devonian land-sea interaction: Evolution of ecosystems and climate (DEVEC). pp. 78-79. [diversity includes 9 ophiuroid spp., 2 asteroid spp.]
- Haude, Reimund and Erich Thomas. 1983. Ophiuren (Echinodermata) des hohen Oberdevons im nördlichen Rheinischen Schiefergebirge.--Paläont. Z. 57(1/2):121-142, Stuttgart. [source Pek & Prokop 1986] [good information/drawings of vertebrae of Furcaster, Ophiaulux decheni, and Onychaster ?velbertensis n. sp.] [Protaster ? ubaghsi n. sp.; juvenile Protasteridae; Stephanoura belgica]
- Haude, Reimund and Erich Thomas. 1989. Ein Oberdevon-/UnterKarbon-Profil im Velberter Sattel (Nördliches Rheinisches Schiefergebirge) mit neun Arte von (?)Sostrocrinus (Echinodermata).--Bulletin de la Société belge de Géologie 98(3/4):373-383. [locality of fossil ophiuroids and asteroids mentioned]
- Haude, Reimund and Erich Thomas. 1994. Eleutherozoen (Echinodermata) aus dem Unter-Karbon von Aprath im Bergischen Land. In C. Hackler, A. Heinrich & E.-B. Krause (eds.), Archäologie im Ruhrgebiet 1994. Geologie, Paläontologie und Vor- und Frühgeschichte Zwischen Lippe und Wupper: 115-132. [includes Chattaster hueffneri, Furcaster sp., Stephanoura? sp.]

July 13, 2010

- Havlicek, Vladimir and Jire Vanek. 1966. The biostratigraphy of the Ordovician of Bohemia. Sbornik Geologických Ved (Paleontologie) rada P. sv 8, pp. 7-70, 16 pls.
- Hawkins, H. L. 19???. William Kingdom Spencer 1878-1955. Biogr. Mem. Roy. Soc. 2: 291-298, portr.
- Hawkins, H. L. and S. M. Hampton. 1927. The occurrence, structure, and affinities of Echinocystis and Palaeodiscus.--Quarterly Journal of the Geological Society of London 83(4):574-603 + pls. 44-46. [includes a stratigraphical account of the Starfish Quarry at Church Hill, Leintwardine (Furcaster, Lapworthura, L. miltoni); a 'starfish bed' with Lapworthura in a quarry at Martin's Shell] [discussion by Gregory, Spencer & Bather on comparing structure of Palaeodiscus with asteroids]
- Hedde, Duncan. 1995. The descent of the Asteroidea and the reaffirmation of paxilloidan primitiveness. Pp. 179-183 in Echinoderm Research 1995, Emson, Smith & Campbell (eds.), Balkema, Rotterdam. [mention of somasteroids, Paleozoic asteroids, Siluraster characteri]
- Herringshaw, Liam. 2000. Strange Silurian starfish. [abstract]. The Palaeontological Association Newsletter No. 45, p. 35. [Lepidaster grayi and four new spp., two of which belong to new genera, in the Much Wenlock Limestone] [suggests that multiradiate state has evolved independently more than once in starfish history]
- Herringshaw, Liam. 2001. Lucky for some – 13-rayed Silurian starfish and the origins of multiradiate asteroids. The Palaeontological Association 45th Annual Meeting Abstracts, p. 18 [in The Palaeontological Association Newsletter No. 48.] [Lepidaster grayi, Much Wenlock Limestone Formation, Dudley, England]
- Herringshaw, L. G., M. P. Smith, and A. T. Thomas. 2007. Evolutionary and ecological significance of Lepidaster grayi, the earliest multiradiate starfish. Zoological Journal of the Linnean Society 150:743-754. [list of Paleozoic multiradiate asteroids; includes Lepidasterella; also a Silurian, Ludlow, Australia, unnamed multiradiate]
- Herringshaw, L. G., A. T. Thomas, and M. P. Smith. 2007. Starfish diversity in the Wenlock of England. Palaeontology 50(5):1211-1229. [Hudsonaster? carectum n. sp.; Doliaster brachyactis n.g. n.sp.; Siluraster? ketlyi, Lepidaster grayi, Lepidactis wenlocki, Palasterina orchilocalia n.sp.] [plus mention of other taxa] [p. 1227 separation of MAPP from IMM frame in Lepidaster provides greater oral flexibility and is absent in earlier asteroids] [study for terminology]
- Hertz, M. 1926. Die Ophiuriden der Deutschen Südpolar-Expedition. -- Deutsche Südpolar-Expedition 19:56pp + 9 plates. [Aganaster gregarius mentioned on p. 54; also Triassic

July 13, 2010

Acrura squamosa; discussion of phylogeny of families of living ophiuroids]

- Hess, Hans. 1955. Die fossilen Astropectiniden (Asteroidea): Neue Beobachtungen und Übersicht über die bekannten Arten. -- Schweizerischen Paläontologischen Abhandlungen 71:1-113 + plates 1-4. [p. 107, fig. 62 Xenaster margaritatus, L. Devonian, p. 107-108]
- Hess, R. 1983. Das Spurenfossil Asteriacites im klastischen Permoskyth (Prebichl-Schichten) der südlichen Admonter Schuppenzone (Ostalpen) und sein paläogeographische Bedeutung. Neues Jahrbuch für Geologie und Paläontologie Monatshefte 1983:513-519. [Wilson & Rigby 2000 suggest latest Permian]
- Hicks, ?. 1873. On the Tremadoc rocks in the neighborhood of St. David's, South Wales, and their fossil contents. Quart. Journ. Geol. Soc. Lond., vol. 29, pp. 39-52, pls. 3, 4.
- Hills, Edwin S. 1929. The geology and palaeontology of the Cathedral Range and the Blue Hills, in North-Western Gippsland. Proceedings of the Royal Society of Victoria, vol. 41 (new series), part II, pp. 176-201, pls. XVII-XVIII, 3 text-figs. [Taeniaster(?) sp. nov. aff. spinus Billings, Yeringian (Upper Silurian) on p. 179.]
- Hisinger, ?. ?????. Lethea Sueca, p. 89, pl. 26, fig. 6. [Source WKS pt. 5, p. 228.]
- Hoare, R. D. and J. Pojeta Jr. 2006. Ordovician polplacophora (Mollusca) from North America. Journal of Paleontology 80(3)supplement: Paleontological Society Memoir 64, i + 27 pp. [p. 26 Stenaster cf. S. obtusus listed as associated fauna, USGS locality 5101-CO, lower Curdsville Limestone; cites Branstrator 1979]
- Hoare, R. D. and M. T. Sturgeon. 1984. Pennsylvanian echinoid pedicellariae. Journal of Paleontology 58(3):895-897. [FH comments that their Fig. 1 items H-L are similar to the adambulacral plates of Umerophiura in Boczarowski 2001:Fig. 3]
- Hohenstein, P. 2004. X-ray imaging for palaeontology. The British Journal of Radiology 77:420-425. [images and practical X-ray methodology] [Fig. 6 of Taeniaster benecke is the specimen with tube feet -- see Glass & Blake 2004]
- Horner, John R. & Richard Lund. 1985. Biotic distribution and diversity in the Bear Gulch Limestone in central Montana. Compte Rendu, Neuvième Congrès Intern. Strat. Géol. Carbonifère, v. 5, pp. 437-442. (IX-ICC v. 5:437-442) [p. 439 fauna includes starfish; mention only]
- Horowitz, A. S. 1991. A fossil site near Sulpher, Indiana (Chesterian, Mississippian). MAPS Digest Expo XIII issue (Mid-America Paleontology Society) pp. 61-70. [faunal list compiled from S. M. Kelly 1984 PhD Thesis; includes Neopalaeaster sp., Tremataster sp.]

July 13, 2010

- Horowitz, A. S. & J. A. Waters. 1972. A Mississippian echinoderm site in Alabama. *Journal of Paleontology* 46(5):660-665. [Monteagle Limestone, Weatherly Mountain, Madison Co. Diverse crinoid/blastoid/other fauna. Includes Onychaster sp. in private collns & in Undiana Univ. collns.]
- Hoskins, Donald M. 1964. Fossil collecting in Pennsylvania. Pennsylvania Geol. Survey 4th ser. Bull. G40 (General Geology Report), 126 pp. [Hallaster at Swatara Gap; surrogate rendition.] [see Hoskins, Inners & Harper 1988]
- Hoskins, Donald M., Jon D. Inners & John A. Harper. 1988. Fossil collecting in Pennsylvania. Pennsylvania Geological Survey, General Geology Report 40, 215 pp. [the 1988 edition is termed the second printing of the third edition of 1983, but actually there are some small differences from the 1983 edition that are mentioned in the preface] [pp. 37-38 Stelleroidea] [Protasterina on p. 96, locality 21 near Roxbury in Franklin County] [Mesopalaeaster and Protasterina on p. 116, locality 29 Swatara Gap in Lebanon County, and p. 198 = plate 2] [what was called Hallaster in the Hoskins 1964 edition is now labeled Protasterina]
- Hotchkiss, F. H. C. 1970. North American Ordovician Ophiuroidea. The genus Taeniaster Billings, 1858 (Protasteridae). *Proc. Biol. Soc. Wash. (D.C.)*, vol. 83, no. 5, pp. 59-76, 10 figs.
- Hotchkiss, F. H. C. 1974. Studies on Paleozoic ophiuroids and the ancestry of the Asterozoa. Ph. D. dissertation, Yale University, June 1974, vi + 147 pp., 24 pls. Abstract in *Dissertation Abstracts International* 35 (6)
- Hotchkiss, F. H. C. 1976. Devonian ophiuroids of New York State. Reclassification of Klasmura, Antiquaster, and Stenaster into the Suborder Scalarina nov., Order Stenurida.--*New York State Museum Bulletin* 425:1-39. [synopsis of NY State Paleozoic ophiuroids] [K. macropleura n.sp.][Mastigactis removed from Encrinasteridae to the Protasteridae]
- Hotchkiss, F. H. C. 1977. Ophiuroid Ophiocanops (Echinodermata) not a living fossil. *J. nat. Hist.* 11: 377-380. [Phragmactis grayae from Swatara Gap; lists Stenaster, Rhopalacoma, Antiquaster, Encrinaster, Taeniaster, Hallaster, Furcaster, Argentinaster.] [Platasterias from Panama, = new locality]
- Hotchkiss, F. H. C. 1978. Studies on echinoderm ray homologies: Lovén's law applies to Paleozoic ophiuroids.--*Journal of Paleontology* 52:537-544. [some Leintwardine specimens of Furcaster leptosoma are reclassified as cf. Loriolaster gracilis Lehmann (new record)] [Encrinaster sp., Devonian, NY, new locality]

July 13, 2010

- Hotchkiss, F. H. C. 1980. The early growth stage of a Devonian ophiuroid and its bearing on echinoderm phylogeny.--*Journal of Natural History* 14:91-96.
- Hotchkiss, F. H. C. 1985. [see Sass & Condrate, 1985.]
- Hotchkiss, F. H. C. 1992. Footnotes on Lovén's law. abstracts for the Second North American Friends of Echinoderms Conference and Workshop, Harbor Branch Oceanographic Institution, Inc., 9-11 July 1992. [distributed at the meeting]
- Hotchkiss, F. H. C. 1993. A new Devonian ophiuroid (Echinodermata: Oegophiurida) from New York State and its bearing on the origin of ophiuroid upper arm plates.--*Proceedings of the Biological Society of Washington*, vol. 106, no. 1, pp. 63-84. [revision of Strataster, S. maciverorum n. sp., S. ohioensis, S. wrighti n. comb., all having upper arm plates; madreporite in interradius III/IV] [revised homologies of plates of ophiuroids, asteroids & crinoids; edrioasteroid ancestry of the Asterozoa is considered] [Oegophiurida rediagnosed]
- Hotchkiss, F. H. C. 1995. Lovén's law and adult ray homologies in echinoids, ophiuroids, edrioasteroids and an ophiocistioid (Echinodermata: Eleutherozoa). *Proceedings of the Biological Society of Washington* [DC], vol. 108, no. 3, pp. 401-435.
- Hotchkiss, F. H. C. 2004. Subphylum Asterozoa species descriptions. In C. E. Brett & F. R. Caudill (eds.) *Geologic overview of the Trenton Group at West Canada Creek, New York*. Department of Invertebrate Paleontology, Museum of Comparative Zoology, Harvard University. [http:// www .mcz .harvard .edu /Departments /InvertPaleo /Trenton /Intro /trentonintro .htm](http://www.mc.z.harvard.edu/Departments/InvertPaleo/Trenton/Intro/trentonintro.htm)
- Hotchkiss, F. H. C., D. K. Armstrong & D. M. Rudkin. 1995. A new occurrence of the genus Euzonosoma (Echinodermata: Ophiuroidea) from the Middle Ordovician Gull River Formation (Caradocian) of Ontario - first North American Ordovician record of the Encrinasteridae. Pp. 483-486 In: J.D. Cooper, M.L. Droser & S.C. Finney (eds.): *Ordovician Odyssey: Short papers for the Seventh International Symposium on the Ordovician System, Las Vegas, Nevada, USA, June 1995*, 498 pp., The Pacific Section Society for Sedimentary Geology (SEPM). Fullerton, California. [Euzonosoma n.sp.; interpreted as robust epifaunal ophiuroid, possibly living oral-side-up; regenerating arm tip] [quarry described by Brett & Rudkin 1997]
- Hotchkiss, F. H. C. and A. M. Clark. 1976. Restriction of the family Poraniidae sensu Spencer & Wright, 1966 (Echinodermata: Asteroidea).--*Bulletin of the British Museum (Natural History) Zoology* 30:263-268 + pls. 1-3.
- Hotchkiss, F. H. C., and A. Glass. 2010. Bdellacoma in the Hunsrück Slate (Lower Devonian):

July 13, 2010

reidentification of Urasterella verruculosa (Asteroidea: Bdellacomidae). pp. 15-21 in Harris et al. (eds.), *Echinoderms: Durham*, Taylor & Francis Group, London. [Proceedings of the 12th International Echinoderms Conference, August 2006, Durham, NH.] [also 5-rayed and multiradiate Bdellacomidae sp. from the Late Devonian Genesee Group, Cortland County, New York (Yale Peabody Museum, det. FH, collected by C.W. Thayer)]

Hotchkiss, F. H. C. and R. Haude. 2004. Observations on Aganaster gregarius and Stephanoura belgica (Ophiuroidea: Ophiolepididae) (Early Carboniferous and Late Devonian age). pp. 425-431 in T. Heinzeller & J. H. Nebelsick (eds.) *Echinoderms: München*. Taylor & Francis Group, London. xxvi + 633 pp. [conference paper presented by R. Haude; see Haude & Hotchkiss 2003]

Hotchkiss, F. H. C., Rudolf J. Prokop and Václav Petr. 1999. Isolated skeletal ossicles of a new brittlestar of the Family Cheiropterasteridae Spencer, 1934 (Echinodermata: Ophiuroidea) in the Lower Devonian of Bohemia (Czech Republic). *Journal of the Czech Geological Society* 44:189-193.

Hotchkiss, F. H. C., Rudolf J. Prokop and Václav Petr. 1999. Isolated vertebrae of brittlestars of the Family Klasmuridae Spencer, 1925 (Echinodermata: Ophiuroidea) in the Devonian of Bohemia (Czech Republic). *Journal of the Czech Geological Society* 44:329-333.

Hotchkiss, F. H. C., Rudolf J. Prokop and Václav Petr. 2007. Isolated ossicles of the Family Eospondylidae Spencer et Wright, 1966, in the Lower Devonian of Bohemia (Czech Republic) and correction of the systematic position of eospondylid brittlestars (Echinodermata: Ophiuroidea: Oegophiurida). *SBORNÍK NÁRODNÍHO MUZEA V PRAZE, Řada B – Přírodní vědy*, 63(1): 3–18; [= *ACTA MUSEI NATIONALIS PRAGAE*, Series B – *Historia Naturalis*, 63(1): 3–18] [Eospondylus is not a euryalid]

Hotchkiss, F. H. C., D. M. Rudkin and S. Anderson. 1997. A case for rearmament- the oldest known evidence of regeneration in sea stars. [abstract] 7th Canadian Paleontology Conference, University of Saskatchewan, Saskatoon, September 26-30, 1997. [see also Rudkin, Hotchkiss & Anderson 1997]

Hotchkiss, F. H. C., David M. Rudkin and Derek K. Armstrong. 1995. Euzonosoma (Echinodermata: Ophiuroidea) from the Middle Ordovician Gull River Formation in Ontario - Some palaeogeographic and palaeoecological implications of the first North American Record of the Encrinasteridae, (Abstract). In: P. A. Johnston (ed.): *Canadian Paleontology Conference, Program and Abstracts No. 5: [The Fifth Canadian Paleontology Conference and International Symposium on the Paleobiology and Evolution of the Bivalvia. Joint Meeting, Royal Tyrrell Museum of Palaeontology, Drumheller, Alberta, 29 September - 2 October 1995.] [CPC-V Drumheller '95]*

July 13, 2010

- Hotchkiss, F. H. C. and P. von Bitter. 1998. Diminutive ophiuroids (Echinodermata) from the upper Windsor Group (Lower Carboniferous: Late Viséan/Asbian) of Nova Scotia, Canada (Abstract). Canadian Paleontology Conference Program and Abstracts No. 8 [The Eighth Canadian Paleontology Conference, 22-25 October 1998][CPC-VIII Collingwood 1998], p. 11. [Also: Royal Ontario Museum Twentieth Annual Research Colloquium, 18 November 1998, Toronto, Abstracts of Papers, p. 8 (abbreviated abstract).]
- Howard, J. D. and I. B. Singh. 1985. Trace fossils in the Mesozoic sediments of Kachchh, Western India. *Palaeogeography, Palaeoclimatology, Palaeoecology* 52:99-122. [source Mangano et al. 2002]
- Hudson, G. H. 1912. A fossil starfish with ambulacral covering plates. *Ottawa Naturalist* vol. 26, pp. 21-26, 45-52, pls. 1-3. [Eostelleroidea n. order, Protopalaeasteridae n. fam., Protopalaeaster narrawayi n.g. n.sp.]
- Hudson, G. H. 1913. The use of the stereogram in paleobiology. *Bull. N. Y. State Mus.*, no. 164, pp. 103-130, pls. 1-13. [Palaeaster parviusculus Billings, holotype, McGill College No. 1586; Protopalaeaster narrawayi Hudson, holotype; Palaeaster niagarensis Hall, Cornell Univ. No. 7331.]
- Hudson, G. H. 1913. Does the type of Protopalaeaster narrawayi present the oral or aboral aspect? *Ottawa Naturalist*, vol. 27, pp. 77-84, pls. 8, 9.
- Hudson, G. H. 1916. On the genus Urasterella with description of a new species. *New York State Museum Bull.* 187, Twelfth report of the Director, 1915, pp. 117-163. [WKS pt. 5 p. 205.] [Urasterella medusa n. sp., Trenton Falls, N.Y., specimen in MCZ; U. pulchella (Billings).]
- Hunter, A. W., B. David and R. Mooi. 2006. Phylogeny of Lower Palaeozoic asterozoans using skeletal homologies derived from the extraxial-axial theory (abstract). 12th International Echinoderm Conference, 7-11 August 2006, University of New Hampshire, Durham, NH. Book of abstracts, p. 34.
- Hunter, A. W., B. Lefebvre, E. Nardin, S. Regnault, P. Van Roy, and S. Zamora. 2006. New echinoderm Lagerstätten from the Upper Ordovician of the eastern Anti-Atlas, Morocco (abstract). 12th International Echinoderm Conference, 7-11 August 2006, University of New Hampshire, Durham, NH. Book of abstracts, pp. 34-35. [protasterid indet, lower Lower Ktaoua Fm, early-middle Caradoc] [eophiurid indet., Lower Ktaoua Fm, earliest Ashgill, starfish bed]
- Hunter, A. W., B. Lefebvre, S. Régault, P. Roussel and R. Claverie. 2006. The oldest recorded mixed ophiuroid-stylophoran assemblage (Echinodermata) from the Middle Ordovician

July 13, 2010

- (Llandeilian) of Gourin (Brittany, France) [abstract]. Journées Georges Ubaghs, 30-31 Janvier 2006, Université de Bourgogne, Dijon (France), Program & Abstracts [B. Lefebvre, B. David, E. Nardin & E. Poty (eds.)] pp. 21-23. [Schistes de Postolonnec Fm] [Taeniaster and Mitrocystella]
- Hunter, A. W., B. Lefebvre, S. Régnault, P. Roussel and R. Claverie. 2007. A mixed ophiuroid -- stylophoran assemblage (Echinodermata) from the Middle Ordovician (Llandeilian) of western Brittany, France. Geological Society, London, Special Publications 275:71-86. [not seen] [Taeniaster armoricanus sp. nov.] [“oldest evidence for gregarious mode of life for ophiuroids”]
- Hunter, A. W. and D. Vizcaïno. 2005. New somasteroids and ophiuroids from the Lower Ordovician of the Montagne Noire (southern France) and their significance to the early evolution of the asterozoans [abstract]. Palaeontological Association Newsletter 60:43. [new specimens and old] [Saint-Chinian Formation (upper Tremadoc-Arenig)] [Landeyran Formation (middle Arenig)] [see Vizcaino & Lefebvre 1999]
- Hunter, A. W. and D. V. Vizcaïno. 2006. New somasteroids and ophiuroids from the Lower Ordovician of the Montagne Noire (southern France) -- their significance to the early evolution of the asterozoans [abstract]. Journées Georges Ubaghs, 30-31 Janvier 2006, Université de Bourgogne, Dijon (France), Program & Abstracts [B. Lefebvre, B. David, E. Nardin & E. Poty (eds.)] pp. 20-21. [New somasteroids and asteroid from the Saint-Chinian Fm] [new ophiuroids from the Landeyran Fm] [the asteroid is cf. hudsonasterids]
- Huxley, T. H. 1878. Catalogue of the Cambrian and Silurian fossils in the Museum of Practical Geology.
- Huxley, T. H. and R. Etheridge. 1865. A catalogue of the collection of fossils in the Museum of Practical Geology, with an explanatory introduction. HMSO, London. lxxix + 381 pp.
- Hyman, L. H. 1955. The invertebrates: Echinodermata. Volume IV. McGraw-Hill, New York, 763 pp.
- Hynda, V. A. and G. M. Pomyanovskaya. 1971. On the Echinodermata from Ordovician Volhynian. Paleontologicheskii Sbornik No. 8, pp. 47-50. [western Volhyn; isolated ossicles (marginals?) of Asteroidea indet p. 49, figs. 3a, 3b.]